

# Working Alliance and Virtual Motor Rehabilitation in Parkinson Patients

Guillermo Palacios-Navarro  
Universidad de Zaragoza  
Calle Ciudad Escolar S/N  
44003, Teruel, SPAIN  
guillermo.palacios@unizar.es

Sergio Albiol-Pérez  
Universidad de Zaragoza  
Calle Ciudad Escolar S/N  
44003, Teruel, SPAIN  
salbiol@unizar.es

José-Antonio Gil-Gómez  
José-Antonio Lozano Quilis,  
Hermenegildo Gil-Gómez  
Universitat Politècnica de València  
C. Vera S/N, 46022, Valencia, SPAIN  
{jgil,jlozano,hgil}@upv.es

## ABSTRACT

Patients that suffer from Parkinson Disease (PD) have different symptoms such as tremors, stiffness and slowness in the execution of first movements and absence of balance control. This paper introduces the concept of working alliance within a virtual rehabilitation system for Parkinson Disease (PD) patients. On one hand, the use of serious games increases the rehabilitation process in PD patients against traditional rehabilitation. On the other hand, this virtual environment encourages the links of confidence between therapist and patient, creating a stronger Working Alliance, which is vital in the patient rehabilitation process. This work in progress has been designed to examine both the working alliance during the virtual rehabilitation process, as well as the results obtained by patients as the clinical study goes on.

## Categories and Subject Descriptors

H5.1. [Multimedia Information Systems]: Artificial, augmented, and virtual realities; H5.2 [Information Interfaces and Presentation]: User Interfaces – *Graphical user interfaces (GUI), Interaction styles, Screen design, User-centered design*. J3 [Computer Applications]: Life and medical Sciences – *Health, Medical information systems*.

## General Terms

Design, Experimentation, Human Factors.

## Keywords

Human Computer Interaction; Virtual Motor Rehabilitation; Therapeutic Alliance; Parkinson Disease.

## 1. INTRODUCTION

Parkinson disease (PD) is a chronic neurodegenerative disease. This manifests itself as a series of debilitating motor and non-motor symptoms having a significant impact on activities of daily living. Nowadays, approximately four million people worldwide suffer from PD, with an average of 1 out of 500 people. It is estimated that over 1,150,000 people aged over 65 in Europe are affected by PD [1].

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. To copy otherwise, or republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee.  
PervasiveHealth 2014, REHAB workshop, May 20–23, 2014, Oldenburg, Germany.  
Copyright 2014.

The main disorders of PD patients are: postural control and more specifically, disorders in the balance, existing in this type of patients a high percentage of risk of falls [2][3], which alter their quality of life. This kind of problems originates a high cost in worldwide health systems and even a high risk of mortality [4].

Besides the problems of postural control, PD patients also present other important deficits associated with their pathology: muscular inflexibility, injuries in visual-spatial tasks, fear to possible falls, loss of confidence, quakes, and mobility reduction to perform all activities in daily life (A-D-L) [5], tremor, bradykinesia, rigidity and postural instability [6]. Consequently, a detriment (physical and cognitive) will be generated [7].

## 2. RELATED WORK

The concept of Virtual Motor Rehabilitation arises from the need to perform a set of tasks by the clinical specialists in their daily occupation. This kind of novel techniques involves a deep change in the process of the traditional motor rehabilitation in such a way that the therapist-patient affinity (Therapeutic Alliance) is clearly reinforced. The importance of it for a successful therapy has been addressed widely [8][9].

The therapeutic Alliance can be defined as the collaborative bond between therapist and patient [10]. According to Bordin [11], the Therapeutic Alliance (TA) consists of three components: the positive bond which develops between a client and a counselor in the counseling process; agreement about the tasks of counseling treatment as well as agreement about the counseling goals. The TA is established through the interpersonal interactions between the counselor and the client in communication process. Nevertheless, this type of interrelation is not curative, but it fosters the compliance and follow-up of the therapeutic process.

On the other hand, Peschken and Johnson [12] stated that the confidence between therapist and patient enhances the degree of empathy, congruence and compliance. In the light of the foregoing, the Working Alliance is a joint construction between patient and therapist, in such a way that opinions and expectations developed all along the process are vital for a suitable establishment of the working alliance [13].

For these reasons, it is suitable to establish the corresponding links between both the therapist and the patient within the process of virtual motor rehabilitation. It is also convenient, before the establishment of the design and work methodology, to create a type of methodology that fulfills this type of interrelationships.

As far as Virtual Motor Rehabilitation is concerned, several sources [14][15] state that patient interest in performing repetitive tasks involved in traditional treatments is an important drawback

in the way of rehabilitation. Indeed, rehabilitation is by its nature repetitive, and repetition tends to “decouple” the mind, and reduce patient’s motivation [16]. Another characteristic deals with the predominance of simple mechanical devices with little or no computerized sensing. Traditional rehabilitation is done one-to-one, meaning one therapist working with one patient. In fact, traditional approaches include exercises often considered repetitive and boring for patients [17]. To overcome this, virtual rehabilitation and serious games can augment physical and cognitive rehabilitation oriented to a significant therapeutic benefit. Virtual rehabilitation can be defined as computer-based technology providing a multisensorial environment with which the user can interact [18].

Technological advances have been gradually incorporated aimed at improvement people quality of living. The application of virtual reality technology for the rehabilitation of cognitive and motor deficits has been growing in the last decade and stroke patients have been one of the main target populations for these new rehabilitation methods [19]. These virtual reality based-methods can offer the patients to be part of immersive experiences that are engaging and rewarding for them.

Some examples can be found in Betker et al. [20], Ma and Bechkoum [21] allowing patients to interact with virtual objects in real-time through multiple modalities and to practice specific motor skills. Conconi et al. [22] introduced a platform for rapid development of serious games, with a special focus on therapeutic support games for behavioral and addictive disorders. Burke et al. [19] developed several games designed for upper limb stroke rehabilitation, which use low-cost webcams as input technology to capture video data of user’s movements.

The advantages associated with the use of Virtual Rehabilitation are numerous. The same hardware can be used for various types of patients (children with attention deficits, post-stroke patients, etc.), as well as for various types of exercises done on those patients. Thus a major advantage in all forms of Virtual Rehabilitation is economy of scale [16]. Another great advantage is interactivity and motivation [23][24] By providing visual and auditory rewards, such as displaying gratifying messages in real time (“Great”, “Very Good”, etc.) or visual stimuli [25], patients are motivated to exercise. Immediate feedback on patient performance and results [26] seems also to be a remarkable help in the rehabilitation process of patients. Games also require cognitive and motor activity so they can engage a person’s attention [15], and most games offer increasingly difficult levels that give the player the sense of challenge in his progress and in such a way that is also adapted to his skills. Another important aspect is that games distract the patient’s attention and can be used to aid in the management of pain [15] [19].

It has been showed that games contribute to increase motivation in rehabilitation sessions, which is the major problem in therapy sessions, caused by the repetitive nature of exercises. Literature is plenty of “serious games” for rehabilitation purposes. This concept has been defined by Michael and Chen [27] as “games that do not have entertainment, enjoyment or fun as their primary purpose”. Serious games are not only the application of games and games technology, they are entertaining, enjoyable and fun, but their main purpose is other that was conceived by the game designer when designing the game or that the user defined when played the game.

Information Technologies (IT) used in the framework of health sciences allow to establish a narrower relation between the patient and the specialist, as well as to increase the communication between both [28]. Within the area of patients' rehabilitation, we find a series of associations and forums involved in facilitating relevant information about rehabilitation processes, where experiences are shared, diagnoses generation done by physicians, questions related to symptoms in specific conditions, answers raised by patients with PD during the rehabilitation process, inherent information in last stages in PD, affective connections among patients having this type of ailments, relatives and physical therapists, forums comprised of medical experts who solve any situation and analyze rehabilitation sessions. Furthermore, we can find communities of scientists, rehabilitators, clinical staff and patients committed to collaboration with those experiences of special relevancy related to this type of ailments.

Our proposal deals with the evaluation of the working alliance within a virtual rehabilitation system (PARK-Rehab System) based on serious games. We think this association will improve the effectiveness of rehabilitation in PD patients.

### **3. METHODS**

#### **3.1 Participants**

We have chosen 4 PD patients belonging to the Teruel PD Patients’ Association. In this experiment, only one group of patients has been considered: the virtual group, which will use our virtual system. The inclusion criteria are: subjects needs to obtain a score in the Mini Mental State Examination greater than 23 (>23), in order to ensure the correct participation; patients should be between 50 and 75. The exclusion criteria are: clinical instability, patients with a history of falls and patient refusal. All the patients have movement slowdown and thinking slowdown. Two of them have rigidity, one has tremor, and one has hearing impairment.

#### **3.2 Instrumentation**

The PARK-Rehab system was designed and developed using UNITY Software and it uses low cost devices such as Tablet PCs. Unity is a development engine of 3D and interactive content that allows publishing/exporting to different platforms, included mobile platforms (iOS, Android). We have chosen this tool taking into account the following reasons: it is free for non-commercial use, it allows the importation of 3D models from the majority of existing platforms and it is relatively simple to use. Three programming languages are available: JavaScript, C# and Boo. We have use C#. The created scripts are added to the 3D models imported to the scene in order to indicate the object behavior as well as its interaction with other objects. To make virtual rehabilitation more feasible for patients, Tablet PCs have been selected, as most of our patients have such devices at home and are familiar with them.

At the moment, two well-known games have been implemented inside the system: ‘Simon’ game and the ‘Dominoes’ game. They both are used by PD Patients in their traditional rehabilitation. Figure 1 shows a PD Patient playing the dominoes game. See Table I for a short description of both games

#### **3.3 Training Programme**

The study was carried out in the Teruel Parkinson Patients Association.



**Figure 1. A PD patient playing the dominoes game.**

The PD patients will perform a total of 20 30-minutes-sessions using PARK-Rehab, twice a week in a pre-fixed timetable.

### 3.4 Metric Analysis

Data gathered from each interaction session is stored in separated files and can be individually processed. The data samples are analyzed to derive a series of metrics which are proposed to indicate the severity of various PD symptoms. Initially, we have considered the following metrics:

The Performance Rate (PR) or (% correct clicks) provides an indication of the impact of PD motor symptoms on the completion of fine motor tasks. This is calculated by comparing the number of correct clicks to the number of incorrect clicks. A correct click is defined as a button press performed while the cursor intersects with the desired target. An incorrect click is defined as any other unnecessary button press.

The Level of Difficulty (LD) provides the maximum level obtained by the patient in every session. It is supposed to be increasing as the number of session increase.





The Completion Time (CT) (milliseconds) describes the total time taken to complete tasks which are not time limited.

For the measurement of the working alliance between the patients and their therapists, the short form of the Working Alliance Inventory (WAI) [29] will be used. The patient/therapist short forms of the WAI comprise 12 items, each four measuring the goal, task and bond aspects of the working alliance. The WAI items were rated separately and independently by the patients and their respective primary therapists on a 7-point Likert scale ranging from 1 = 'not at all' to 7 = 'a lot'. Prior to the completion of the questionnaires, both the patients and therapists will be informed that their ratings would be treated as confidential. The questionnaire regarding their working alliance will be performed at the end of session 2 and again at the end of session number 20 by both therapists and patients.

## 4. DISCUSSION AND CONCLUSIONS

The purpose of this work in progress deals with the examination of development and interaction of the working alliance during the process of virtual rehabilitation in PD patients. Furthermore, a series of metrics are stored for every patient, in order to show their evolution along time.

**Table 1. Implemented games in Park-Rehab System**

DOMINOES GAME	
Virtual Game	Visual Aspect
A series of dominoes pieces are shown to the patient. A score showing the hits, errors and remaining pieces to complete the task	
The patient select the the dominoes pieces whose sum is equal or greater than six.	
SIMON GAME	
A Control panel showed in the initial screen allows the therapist to enter the patient's data.	
The patient has to remember the presented sequence by pushing the right buttons. The level of difficulty is more and more challenged	

At the moment we are immersed in the virtual rehabilitation process, and first inputs coming from both therapists and PD patients are promising, showing a great deal of commitment in the virtual rehabilitation process. Future work deals with the establishment of two groups of patients: a control group and a virtual group, respectively. In this way, we will be able to show the benefits of virtual rehabilitation in conjunction with the working alliance in the process of PD patients' rehabilitation.

## 5. ACKNOWLEDGMENTS

We want to thank both therapists and patients belonging to the Teruel Parkinson Disease Association, who willingly have collaborated in this project.

This contribution was partially funded by the Gobierno de Aragón, Departamento de Industria e Innovación, y Fondo Social Europeo "Construyendo Europa desde Aragón"..

## 6. REFERENCES

- [1] Andlin-Sobocki, P., Jönsson, B., Wittchen, H. U. and Olesen, J. 2005. Cost of disorders of the brain in Europe. *European Journal of Neurology*, 12, 1–27.
- [2] Bloem, B. R., Van Vugt, J. P. and Beckley, D. J. 2001. Postural instability and falls in Parkinson's disease. *Adv Neurol*, 87, 209–23.
- [3] Franchignoni, F., Martignoni, E., Ferriero, G., Pasetti, C. 2005. Balance and fear of falling in Parkinson's disease. *Parkinsonism Relat Disord*, 11, 427–33.
- [4] Hely, M. A., Morris, J. G., Traficante, R., Reid, W.G., O'Sullivan, D. J., Williamson, P. M. 1999. The sydney multicentre study of Parkinson's disease: progression and mortality at 10 years. *J Neurol Neurosurg Psychiatr*, 67, 300–307.
- [5] Michalowska, M., Fiszer, U., Krygowska-Wajs, A., Owczarek, K. 2005. Falls in Parkinson's disease. Causes and impact of patients' quality of life. *Funct Neurology*, 20 (4), 163–8.
- [6] Synnott, J. WiiPD-Objective Home Assessment of Parkinson's Disease using the Nintendo Wii Remote," unpublished.
- [7] Carne, W. et al. 2005. Efficacy of multidisciplinary treatment program on long-term outcomes of individuals with Parkinson's disease. *J. Rehabil Res Dev*, 42,779–86.
- [8] Lambert, M. J., Barley, D. E. 2002. Research summary on the therapeutic relationship and psychotherapy outcome. In: Norcross JC, editor. *Psychotherapy relationships that work: Therapist contributions and responsiveness to patients*. New York: Oxford University Press, 17–32.
- [9] Constantino, M. J., Castonguay, L. G., Schut. A. J. 2002. The working alliance: A flagship for the 'scientist-practitioner' model in psychotherapy. In: Tryon GS, editor. *The working alliance: A flagship for the scientist-practitioner model in psychotherapy*. Boston, MA: Allyn & Bacon, 81–131.
- [10] Krupnick, J. L. 2006. The Role of the Therapeutic Alliance in Psychotherapy and Pharmacotherapy Outcome: Findings in the National Institute of Mental Health Treatment of Depression Collaborative Research Program. *The Journal of Lifelong Learning in Psychiatry*, IV, 2, 269–277.
- [11] Bordin, E. S. 1979. The generalizability of the psychoanalytic concept of the working alliance", *Psychotherapy: Theory, Research and Practice*, 16(3), 252–260.
- [12] Peschken, W. E. and Johnson, M. E. 1997. Therapist and client trust in the therapeutic relationship. *Psychotherapy Research*, 7(4), 439–447.
- [13] Corbella, S., Botella, L. 2003. La alianza terapéutica: historia, investigación y evaluación. *Anales de Psicología*, 19(2), 205–221.
- [14] Lozano-Quilis, J., Gil-Gómez, H., Gil-Gómez, J., Albiol-Pérez, S., Palacios, G., Fardoun, H. M. and Mashat, A. S. 2013. Virtual reality system for multiple sclerosis rehabilitation using kinect. *Pervasive Computing Technologies for Healthcare (PervasiveHealth)*, 366–369.
- [15] Fardoun H.M., Abdulraham H.A., Paules Cipres A., Ramírez Castillo J., Albiol-Pérez S. 2013. CRehab: A Cloud-based Framework for the Management of Rehabilitation Processes. *Pervasive Computing Technologies for Healthcare (PervasiveHealth '13)*, 397–400, 2013.
- [16] Burdea, G. C. 2003. Virtual rehabilitation-benefits and challenges. *Methods Inf Med*, 42 (5), 519–523.
- [17] Merola, A. et al. 2011. Parkinson's disease progression at 30 years: a study of subthalamic deep brain-stimulated patients. *Brain*, 134 (7), 2074–2084.
- [18] Levin, L. 2011. Can virtual reality offer enriched environments for rehabilitation. *Exp Rev Neurotherapeut*, 11,153–155.
- [19] Burke, J. W. et al. 2009. Optimising engagement for stroke rehabilitation using serious games. *Visual Computer*, 25, 1085—1099.
- [20] Betker, A. L., Desai, A., Nett, C., Kapadia, N. and Szturm T. 2007. Gamebased Exercises for Dynamic Short-Sitting Balance Rehabilitation of People with Chronic Spinal Cord and Traumatic Brain Injuries. *Physical Therapy*, 87, 10, 1389—1398.
- [21] Ma, M., Bechkoum, K. 2008. Serious Games for Movement Therapy after Stroke. In *Proceedings of the IEEE Int. Conf. on Systems, Man and Cybernetics*, 1872—1877.
- [22] Conconi, A., Ganchev, T., Kocsis, O., Papadopoulos, G., Fernández-Aranda, F., Jiménez-Murcia, S. 2008. PlayMancer: A Serious Gaming 3D Environment. In *Proceedings of the Int. Conf. on Automated Solutions for Cross Media Content and Multi-channel Distribution. AXMEDIS'08*, IEEE Press, 111—117.
- [23] Riess, T. J. 1998. Gait and Parkinson's disease: a conceptual model for an augmented-reality based therapeutic device. *Stud Health Technol Inform*, 58, 200–208.
- [24] Mirelman, A., Maidan, I., Herman, T., Deutsch, J. E., Giladi, N., Hausdorff, J. M. 2011. Virtual reality for gait training: can it induce motor learning to enhance complex walking and reduce fall risk in patients with Parkinson's disease?. *J Gerontol A Biol Sci Med Sci*, 66, 234–40.
- [25] Deutsch, J., Borbely, M., Filler, J., Huhn, K., Guarrera-Bowlby, P. 2008. Use of a low-cost, commercially available gaming console (Wii) for rehabilitation of an adolescent with cerebral palsy. *Phys Ther*, 88, 1196–207.
- [26] Fong, K. N. K. et al. 2010. Usability of a virtual reality environment simulating an automated teller machine for assessing and training persons with acquired brain injury. *J Neuroeng Rehabil*, 7:19.
- [27] Chen, S., Michael, D. 2006. *Serious Games: Games that Educate, Train and Inform*. Boston: Thomson Course Technology PTR.
- [28] Kruse, R.L. et al. 2012. Internet Use by Primary Care Patients: Where Is the Digital Divide?. *Family Medicine*, 44 (5)
- [29] Tracey, T. J., Kokotovic, A. M. 1989. Factor structure of the working alliance inventory. *Psychological Assessment: A Journal of Consulting and Clinical Psychology*, 1: 207–210.